

AMENDMENTS TO THE CLAIMS:


This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1 - 75. (canceled)

76. (new) A method of increasing the vigor and/or the yield of an agronomic plant comprising treating the seed with a composition which comprises an effective amount of a fungicide which has no significant activity against fungal plant pathogens for such agronomic plant and wherein the seed is treated with an inoculant comprising *Azospirillum spp.*, or *Rhizobium spp.*, or *Bradyrhizobium spp.*, or a mixture of *Rhizobium spp.* and *Bradyrhizobium spp.*, or a mixture of either *Rhizobium spp.*, or *Bradyrhizobium spp.* with any other microorganisms.

77. (new) The method according to claim 76, wherein the fungicide comprises a compound having the formula



wherein Z₁ and Z₂ are C or N and are part of an aromatic ring selected from benzene, pyridine, thiophene, furan, pyrrole, pyrazole, thiazole, benzothiophene and isothiazole;

A is selected from --C(X)-amine, --C(O)--SR₃, --NH--C(X)R₄, and --C(=NR₃)--XR₇;

B is --W_m--Q(R₂)₃ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R₄;

Q is C, Si, Ge, or Sn;

W is $--C(R_3)_p H_{(2-p)}--$; or when Q is C, W is selected from $--C(R_3)_p H_{(2-p)}--$, $--N(R_3)_m H_{(1-m)}--$, $--S(O)_p--$, and $--O--$;

X is O or S;

n is 0, 1, 2, or 3;

m is 0 or 1;

p is 0, 1, or 2;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;

b) C₁-C₄ alkyl, alkenyl, alkynyl, C₃-C₆ cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C₁-C₄ alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxycarbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C₁-C₄ alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C₁-C₄ alkoxy, alkenoxy, alkynoxy, C₃-C₆ cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo;

wherein two R groups may be combined to form a fused ring;

each R₂ is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R₄ or halogen; and wherein, when Q is C, R₂ may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino;

wherein two R₂ groups may be combined to form a cyclo group with Q;

R₃ is C₁-C₄ alkyl;

R₄ is C₁-C₄ alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino;

R₇ is C₁-C₄ alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R₄; or an agronomic salt thereof.

78. (new) The method according to claim 77, wherein

Z_1 and Z_2 are C and are part of an aromatic ring which is thiophene;

A is selected from --C(X)-amine, --C(O)—SR₃, --NH--C(X)R₄, and --C(=NR₃)--XR₇ ;

B is --W_m --Q(R₂)₃ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R₄ ;

Q is C, Si, Ge, or Sn;

W is --C(R₃)_p H_(2-p) --; or when Q is C, W is selected from --C(R₃)_p H_(2-p) --, --N(R₃)_m H_(1-m) --, --S(O)_p --, and --O--;

X is O or S;

n is 0, 1, 2, or 3;

m is 0 or 1;

p is 0, 1, or 2;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;

b) C₁-C₄ alkyl, alkenyl, alkynyl, C₃-C₆ cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C₁-C₄ alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxycarbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C₁-C₄ alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C₁-C₄ alkoxy, alkenoxy, alkynoxy, C₃-C₆ cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo;

each R₂ is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R₄ or halogen; and wherein, when Q is C, R₂ may also be selected from halo, alkoxy, alkylthio, alkylamino, and

dialkylamino, and further when Q is C, R₂ may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino; and further when Q is C, then two R₂ groups may be combined to form a cycloalkyl group with Q;

R₃ is C₁-C₄ alkyl;

R₄ is C₁-C₄ alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino; R₇ is C₁-C₄ alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R₄; or an agronomic salt thereof.

79. (new) The method according to claim 77, wherein

Z₁ and Z₂ are C and are part of an aromatic ring which is thiophene;

A is selected from --C(X)-amine, wherein the amine is substituted with a first and a second amine substituent or with an alkylaminocarbonyl and a hydrogen, --C(O)--SR₃, --NH--C(X)R₄, and --C(=NR₃)-XR₇;

the first amine substituent is selected from the group consisting of C₁ - C₁₀ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof optionally substituted with one or more halogen, hydroxy, alkoxy, alkylthio, nitrile, alkylsulfonate, haloalkylsulfonate, phenyl, C₃ - C₆ cycloalkyl and C₅ - C₆ cycloalkylkenyl; phenyl optionally substituted with one or more C₁ - C₄ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof, cycloalkyl, cycloalkenyl, haloalkyl, alkoxy and nitro; C₃ - C₆ cycloalkyl, C₅ - C₆ cycloalkenyl, alkoxy, alkenoxy, alkynoxy, dialkylamino, and alkylthio;

and the second amine substituent is selected from the group consisting of hydrogen; C₁ - C₆ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof optionally substituted with one or more halogen, hydroxy, alkylcarbonyl, haloalkylcarbonyl, alkoxy, alkenoxy, alkynoxy, dialkylphosphonyl;

B is --W_m--Q(R₂)₃ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R₄;

Q is C, Si, Ge, or Sn;

W is --C(R₃)_p H_(2-p)--; or when Q is C, W is selected from --C(R₃)_p H_(2-p)--, --N(R₃)_m H_(1-m)--, --S(O)_p--, and --O--;

X is O or S;

n is 2;

m is 0 or 1;

p is 0, 1, or 2;

wherein two R groups are combined to form a nonheterocyclic ring fused with the thiophene ring, which is not a benzothiophene other than a tetrahydrobenzothiophene, said two R groups being selected from the group consisting of C₁ - C₄ alkyl, alkenyl, C₃ - C₆ cycloalkyl and cycloalkenyl, each optionally substituted with hydroxy, thio, phenyl, C₁ - C₄ alkoxy, alkylthio, alkylsulfinyl, or alkylsulfonyl;

each R₂ is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R₄ or halogen; and wherein when Q is C, R₂ may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino; and further when Q is C, then two R₂ groups may be combined to form a cycloalkyl group with Q;

R₃ is C₁-C₄ alkyl;

R₄ is C₁-C₄ alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino; and

R₇ is C₁-C₄ alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R₄; or an agronomic salt thereof

80. (new) The method according to claim 77, wherein

Z₁ and Z₂ are C and are part of an aromatic ring which is thiophene;

A is --C(X)-amine wherein the amine is an N-bonded heterocyclic compound chosen from the group consisting of morpholine, piperazine, piperidine, and pyrrolidine, each optionally substituted with C₃ - C₆ alkyl groups;

B is --W_m --Q(R₂)₃ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R₄ ;

Q is C or Si;

W is --C(R₃)_p H_(2-p) --; or when Q is C, W is selected from --C(R₃)_p H_(2-p) --, --N(R₃)_m H_(1-m) --, --S(O)_p --, and --O--;

X is O;

n is 2;

m is 0 or 1;

p is 0, 1, or 2;

wherein the two R groups are alkenyl groups and are combined to form a fused ring with the thiophene ring with is benzothiophene; wherein the alkenyl groups are optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C₂ - C₄ alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxycarbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

each R₂ is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, and phenyl, each optionally substituted with R₄ or halogen; and wherein when Q is C, R₂ may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino; or wherein two R₂ groups may be combined to form a cyclo group with Q;

R₃ is C₁-C₄ alkyl; and

R₄ is C₁-C₄ alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino; or an agronomic salt thereof

81. (new) The method according to claim 77, wherein

Z₁ and Z₂ are C and are part of an aromatic ring which is benzothiophene; and

A is selected from --C(X)-amine wherein the amine is an unsubstituted, monosubstituted or disubstituted nonheterocyclic amino radical, --C(O)—SR₃, --NH--C(X)R₄, and --C(=NR₃)--XR₇ ;

B is --W_m --Q(R₂)₃ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R₄ ;

Q is C, Si, Ge, or Sn;

W is --C(R₃)_p H_(2-p) --; or when Q is C, W is selected from --C(R₃)_p H_(2-p) --, --N(R₃)_m H_(1-m) --, --S(O)_p --, and --O--;

X is O or S;

n is 0, 1, 2, or 3;

m is 0 or 1;

p is 0, 1, or 2;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;

b) C₁-C₄ alkyl, alkenyl, alkynyl, C₃-C₆ cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C₁-C₄ alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxycarbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C₁-C₄ alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C₁-C₄ alkoxy, alkenoxy, alkynoxy, C₃-C₆ cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo;

each R₂ is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R₄ or halogen; and wherein, when Q is C, R₂ may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino;

wherein two R₂ groups may be combined to form a cyclo group with Q which is 1-methylcyclopropyl, 1-methylcyclopentyl, or 1-methylcyclohexyl;

R₃ is C₁-C₄ alkyl;

R₄ is C₁-C₄ alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino; and

R₇ is C₁-C₄ alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R₄; or an agronomic salt thereof

82. (new) The method according to claim 77, wherein

Z₁ and Z₂ are C or N and are part of an aromatic ring which is furan; and

A is selected from --C(X)-amine wherein the amine is substituted with a first and a second amine substituent or with an alkylaminocarbonyl and a hydrogen, --C(O)—SR₃, --NH--C(X)R₄, and --C(=NR₃)--XR₇;

the first amine substituent is selected from the group consisting of C₁ - C₁₀ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof optionally substituted with one or more halogen, hydroxy, alkoxy, alkylthio, nitrile, alkylsulfonate, haloalkylsulfonate, phenyl, a 5-membered heteroaryl, C₃ - C₆ cycloalkyl and C₅ - C₆

cycloalkylkenyl; phenyl optionally substituted with one or more C₁ - C₄ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof, cycloalkyl, cycloalkenyl, haloalkyl, alkoxy and nitro; C₃ - C₆ cycloalkyl, C₅ - C₆ cycloalkenyl, alkoxy, alkenoxy, alkynoxy, dialkylamino, and alkylthio;

and the second amine substituent is selected from the group consisting of hydrogen; C₁ - C₆ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof optionally substituted with one or more halogen, hydroxy, alkylcarbonyl, haloalkylcarbonyl, alkoxy carbonyl, and dialkylphosphonyl;

B is --W_m --Q(R₂)₃ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R₄ ;

Q is C, Si, Ge, or Sn;

W is --C(R₃)_p H_(2-p) --; or when Q is C, W is selected from --C(R₃)_p H_(2-p) --, --N(R₃)_m H_(1-m) --, --S(O)_p --, and --O--;

X is O or S;

n is 0, 1, or 2;

m is 0 or 1;

p is 0, 1, or 2;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;

b) C₁-C₄ alkyl, alkenyl, alkynyl, C₃-C₆ cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C₁-C₄ alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxy carbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C₁-C₄ alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C₁-C₄ alkoxy, alkenoxy, alkynoxy, C₃-C₆ cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl,

alkylcarbonyloxy, alkoxycarbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo;

wherein two R groups may be combined to form a fused ring;

each R₂ is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R₄ or halogen; and wherein, when Q is C, R₂ may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino;

wherein two R₂ groups may be combined to form a cyclo group with Q which is 1-methylcyclopropyl, 1-methylcyclopentyl, or 1-methylcyclohexyl;

R₃ is C₁-C₄ alkyl;

R₄ is C₁-C₄ alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino; and

R₇ is C₁-C₄ alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R₄; or an agronomic salt thereof

83. (new) The method according to claim 77, wherein

Z₁ and Z₂ are C and are part of an aromatic ring which is benzothiophene; and

A is selected from --C(X)-amine wherein the amine is an unsubstituted, monosubstituted or disubstituted nonheterocyclic amino radical, --C(O)—SR₃, --NH--C(X)R₄, and --C(=NR₃)--XR₇ ;

B is --W_m --Q(R₂)₃ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R₄ ;

Q is C, Si, Ge, or Sn;

W is --C(R₃)_p H_(2-p) --; or when Q is C, W is selected from --C(R₃)_p H_(2-p) --, --N(R₃)_m H_(1-m) --, --S(O)_p --, and --O--;

X is O or S;

n is 0, 1, 2, or 3;

m is 0 or 1;

p is 0, 1, or 2;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;

b) C₁-C₄ alkyl, alkenyl, alkynyl, C₃-C₆ cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C₁-C₄ alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxycarbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C₁-C₄ alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C₁-C₄ alkoxy, alkenoxy, alkynoxy, C₃-C₆ cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo;

each R₂ is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R₄ or halogen; and wherein, when Q is C, R₂ may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino;

wherein two R₂ groups may be combined to form a cyclo group with Q which is 1-methylcyclopropyl, 1-methylcyclopentyl, or 1-methylcyclohexyl;

R₃ is C₁-C₄ alkyl;

R₄ is C₁-C₄ alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino; and

R₇ is C₁-C₄ alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R₄; or an agronomic salt thereof.

84. (new) The method according to claim 77, wherein

Z₁ and Z₂ are C and are part of an aromatic ring which is furan; and

A is selected from --C(X)-amine wherein the amine is substituted with a first and a second amine substituent or with an alkylaminocarbonyl and a hydrogen, --C(O)—SR₃, --NH--C(X)R₄, and --C(=NR₃)--XR₇;

the first amine substituent is selected from the group consisting of C₁ - C₁₀ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof optionally substituted with one or more halogen, hydroxy, alkoxy, alkylthio, nitrile, alkylsulfonate, haloalkylsulfonate, phenyl, a 5-membered heteroaryl, C₃ - C₆ cycloalkyl and C₅ - C₆

cycloalkylkenyl; phenyl optionally substituted with one or more C₁ - C₄ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof, cycloalkyl, cycloalkenyl, haloalkyl, alkoxy and nitro; C₃ - C₆ cycloalkyl, C₅ - C₆ cycloalkenyl, alkoxy, alkenoxy, alkynoxy, dialkylamino, and alkylthio;

and the second amine substituent is selected from the group consisting of hydrogen; C₁ - C₆ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof optionally substituted with one or more halogen, hydroxy, alkylcarbonyl, haloalkylcarbonyl, alkoxycarbonyl, and dialkylphosphonyl;

B is --W_m --Q(R₂)₃ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R₄ ;

Q is C, Si, Ge, or Sn;

W is --C(R₃)_p H_(2-p) --; or when Q is C, W is selected from --C(R₃)_p H_(2-p) --, --N(R₃)_m H_(1-m) --, --S(O)_p --, and --O--;

X is O or S;

n is 0, 1, or 2;

m is 0 or 1;

p is 0, 1, or 2;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;

b) C₁-C₄ alkyl, alkenyl, alkynyl, C₃-C₆ cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C₁-C₄ alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxycarbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C₁-C₄ alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C₁-C₄ alkoxy, alkenoxy, alkynoxy, C₃-C₆ cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl,

alkylcarbonyloxy, alkoxycarbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo;

wherein two R groups may be combined to form a fused ring;

each R₂ is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R₄ or halogen; and wherein, when Q is C, R₂ may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino;

wherein two R₂ groups may be combined to form a cyclo group with Q which is 1-methylcyclopropyl, 1-methylcyclopentyl, or 1-methylcyclohexyl;

R₃ is C₁-C₄ alkyl;

R₄ is C₁-C₄ alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino; and

R₇ is C₁-C₄ alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R₄; or an agronomic salt thereof.

85. (new) The method according to claim 77, wherein

Z₁ and Z₂ are C and are part of an aromatic ring which is pyridine; and

A is selected from the group consisting of --C(O)—SR₃, --NH--C(X)R₄, and --C(=NR₃)--XR₇ and --C(X)-amine wherein the amine is substituted with alkylaminocarbonyl and a hydrogen or wherein the amine has a first and a second amine substituent;

the first amine substituent is selected from the group consisting of C₁ - C₁₀ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof optionally substituted with one or more halogen, hydroxy, alkoxy, alkylthio, nitrile, alkylsulfonate, haloalkylsulfonate, phenyl, a 5-membered heteroaryl, C₃ - C₆ cycloalkyl and C₅ - C₆ cycloalkylkenyl; phenyl optionally substituted with one or more C₁ - C₄ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof, cycloalkyl, cycloalkenyl, haloalkyl, alkoxy and nitro; C₃ - C₆ cycloalkyl, C₅ - C₆ cycloalkenyl, alkoxy, alkenoxy, alkynoxy, dialkylamino, and alkylthio;

and the second amine substituent is selected from the group consisting of hydrogen; C₁ - C₆ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof optionally substituted with one or more halogen, hydroxy, alkylcarbonyl, haloalkylcarbonyl, alkoxycarbonyl, and dialkylphosphonyl;

B is $--W_m--Q(R_2)_3$ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R_4 ;

Q is C, Si, Ge, or Sn;

W is $--C(R_3)_p H_{(2-p)}--$; or when Q is C, W is selected from $--C(R_3)_p H_{(2-p)}--$, $--N(R_3)_m H_{(1-m)}--$, $--S(O)_p--$, and $--O--$;

X is O or S;

n is 0, 1, or 2;

m is 0 or 1;

p is 0, 1, or 2;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;

b) C_1-C_4 alkyl, alkenyl, alkynyl, C_3-C_6 cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C_1-C_4 alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxycarbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C_1-C_4 alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C_1-C_4 alkoxy, alkenoxy, alkynoxy, C_3-C_6 cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo;

each R_2 is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R_4 or halogen; and wherein, when Q is C, R_2 may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino; or wherein two R_2 groups may be combined to form a cyclo group with Q which is 1-methylcyclopropyl, 1-methylcyclopentyl, or 1-methylcyclohexyl;

R_3 is C_1-C_4 alkyl;

R_4 is C_1-C_4 alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino; and

R₇ is C₁-C₄ alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R₄; or an agronomic salt thereof.

86. (new) The method according to claim 77, wherein

Z₁ and Z₂ are C and are part of an aromatic ring which is benzene; and

A is selected from the group consisting of --C(X)-amine wherein the amine is substituted with a first and a second amine substituent or with an alkylaminocarbonyl and a hydrogen; --C(O)—SR₃, --NH--C(X)R₄, and --C(=NR₃)--XR₇;

the first amine substituent is selected from the group consisting of C₁ - C₁₀ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof optionally substituted with one or more halogen, hydroxy, alkoxy, alkylthio, nitrile, alkylsulfonate, haloalkylsulfonate, phenyl, C₃ - C₆ cycloalkyl and C₅ - C₆ cycloalkylkenyl; phenyl optionally substituted with one or more C₁ - C₄ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof, cycloalkyl, cycloalkenyl, haloalkyl, alkoxy and nitro; C₃ - C₆ cycloalkyl, C₅ - C₆ cycloalkenyl, alkoxy, alkenoxy, alkynoxy, dialkylamino, and alkylthio;

and the second amine substituent is selected from the group consisting of hydrogen; C₁ - C₆ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof optionally substituted with one or more halogen, hydroxy, alkylcarbonyl, haloalkylcarbonyl, alkoxycarbonyl, and dialkylphosphonyl;

B is --W_m --Q(R₂)₃ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R₄ ;

Q is Si, Ge, or Sn;

W is --C(R₃)_p H_(2-p) --;

X is O or S;

n is 0, 1, 2 or 3;

m is 0 or 1;

p is 0, 1, or 2;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;

b) C₁-C₄ alkyl, alkenyl, alkynyl, C₃-C₆ cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C₁-C₄ alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxycarbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C₁-C₄ alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C₁-C₄ alkoxy, alkenoxy, alkynoxy, C₃-C₆ cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo;

each R₂ is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R₄ or halogen;

R₃ is C₁-C₄ alkyl;

R₄ is C₁-C₄ alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino; and

R₇ is C₁-C₄ alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R₄; or an agronomic salt thereof.

87. (new) The method according to claim 77, wherein

Z₁ and Z₂ are C and are part of an aromatic ring which is furan; and

A is selected from --C(X)-amine wherein the amine is substituted with a first and a second amine substituent or with an alkylaminocarbonyl and a hydrogen, --C(O)—SR₃, --NH--C(X)R₄, and --C(=NR₃)--XR₇;

the first amine substituent is selected from the group consisting of C₁ - C₁₀ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof optionally substituted with one or more halogen, hydroxy, alkoxy, alkylthio, nitrile, alkylsulfonate, haloalkylsulfonate, phenyl, a 5-membered heteroaryl, C₃ - C₆ cycloalkyl and C₅ - C₆ cycloalkylkenyl; phenyl optionally substituted with one or more C₁ - C₄ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof, cycloalkyl, cycloalkenyl, haloalkyl, alkoxy and nitro; C₃ - C₆ cycloalkyl, C₅ - C₆ cycloalkenyl, alkoxy, alkenoxy, alkynoxy, dialkylamino, and alkylthio;

and the second amine substituent is selected from the group consisting of hydrogen; C₁ - C₆ straight or branched alkyl, alkenyl, or alkynyl groups or mixtures thereof optionally substituted with one or more halogen, hydroxy, alkylcarbonyl, haloalkylcarbonyl, alkoxy carbonyl, and dialkylphosphonyl;

B is --W_m--Q(R₂)₃ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R₄ ;

Q is C, Si, Ge, or Sn;

W is --C(R₃)_p H_(2-p) --; or when Q is C, W is selected from --C(R₃)_p H_(2-p) --, --N(R₃)_m H_(1-m) --, --S(O)_p --, and --O--;

X is O or S;

n is 2;

m is 0 or 1;

p is 0, 1, or 2;

wherein the two R groups are combined to form a nonheterocyclic ring fused to said furan ring which is not benzofuran when A is --C(X)--amine, B is --W_m(Q)--(R₂)₃, and Q is C or Si, said R groups being selected from the group consisting of C₁ - C₄ alkyl, alkenyl, C₃ - C₆ cycloalkyl and cycloalkenyl, each optionally substituted with hydroxy, thio, phenyl, C₁ - C₄ alkoxy, alkylthio, alkylsulfinyl, or alkylsulfonyl; and

each R₂ is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R₄ or halogen; and wherein, when Q is C, R₂ may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino; wherein further when Q is C, then two R₂ groups may be combined to form a cyclo group with Q;

R₃ is C₁-C₄ alkyl;

R₄ is C₁-C₄ alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino; and


R₇ is C₁-C₄ alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R₄; or an agronomic salt thereof.

88. (new) The method according to claim 77, wherein Z₁ and Z₂ are C and are part of a thiophene ring.


89. (new) The method according to claim 88, wherein A is -C(O)-amine, wherein the amino radical is substituted with one or two groups selected from hydrogen;

hydroxy; alkyl, alkenyl, and alkynyl, which may be straight or branched chain or cyclic; alkoxyalkyl; haloalkyl; hydroxyalkyl; alkylthio; alkylthioalkyl; alkylcarbonyl; alkoxyalkyl; aminocarbonyl; alkylaminocarbonyl; cyanoalkyl; mono-or dialkylamino; phenyl, phenylalkyl or phenylalkenyl, each optionally substituted with one or more C₁ - C₄ alkyl, alkoxy, haloalkyl, C₃ - C₆ cycloalkyl, halo, or nitro groups; and C₁ - C₄ alkyl or alkenyl substituted with pyrimidinyl, thienyl, or furanyl; and wherein the amino radical may be a N-bonded heterocycle selected from morpholine, piperazine, piperidine, pyrrole, pyrrolidine, imidazole, and triazoles, each optionally substituted with C₁ - C₆ alkyl groups.


90. (new) The method according to claim 89, wherein in -W_m, m is 0.
91. (new) The method according to claim 89, wherein Q is Si.
92. (new) The method according to claim 91, wherein each R₂ is C₁ - C₄ alkyl or haloalkyl.
93. (new) The method according to claim 92, wherein each R₂ is methyl.
94. (new) The method according to claim 93, wherein A is alkylaminocarbonyl or dialkylaminocarbonyl.
95. (new) The method according to claim 76, wherein the fungicide is 4,5-dimethyl-*N*-2-propenyl-2-(trimethylsilyl)-3-thiophenecarboxamide.
96. (new) The method according to claim 76, wherein the fungicide comprises a compound having the formula:




(a)




(b)




(c)




(d)




(e)




(f)




(g)



(h)



(i)



(j)

where A is --C(X)-amine; B is --W_m --Q(R₂)₃; and A can be B when B is A except when the formula is f), then Q cannot be Si;

Q is C or Si;

W is --NH--, --O-- or NCH₃ --;

X is O or S;

m is 0 or 1, provided that m is 0 when Q is Si;

n is 0, 1, 2, or 3;

p is 0, 1 or 2, and n plus p is equal to or less than 3;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;


b) C₁–C₄ alkyl, alkenyl, alkynyl, C₃–C₆ cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C₁–C₄ alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxy carbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C₁–C₄ alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C₁–C₄ alkoxy, alkenoxy, alkynoxy, C₃–C₆ cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl, alkylcarbonyloxy, alkoxy carbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo; each R₂ is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R₄ or halogen; and wherein, when Q is C, R₂ may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino; wherein two R₂ groups may be combined to form a cyclo group with Q; R₄ is C₁–C₄ alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino;

or an agronomic salt thereof.

97. (new) The method according to claim 76, wherein the fungicide comprises a compound having the formula:



wherein R^2 is ethyl, iso-propyl, propyl or allyl;

A is $N(CH_3)_{1-n} H_n R^5$ or OR^6 wherein n is 0 or 1, R^5 is $(CH_3)_m (CH_3 CH_2)_{3-m} C$, 1-methyl-1-cyclopentyl, 1-methyl-1-cyclohexyl or 2,3-dimethyl-2-butyl wherein m is 0, 1, 2 or 3 and R^6 is independently R^5 , or 2,3,3-trimethyl-2-butyl;

R^3 is H or independently R^4 ; and

R^4 is halo or CH_3 ;

with the proviso that when A is $N(CH_3)_{1-n} H_n R^5$, if R^3 is H and R^5 is 1-methyl-1-cyclohexyl or $(CH_3)_m (CH_2 CH_3)_{3-m} C$, where m is 0 or 3, or if R^3 is halo and R^2 is $(CH_3)_m (CH_3 CH_2)_{3-m} C$, where m is 3, then R^2 cannot be ethyl;

and with the proviso that when A is OR^6 then m is equal to or less than 2, and if R^3 is H or halo and R^2 is ethyl or isopropyl, then R^6 is $(CH_3)_M (CH_3 CH_2)_{3-M} C$ where m is 1;

or an agronomic salt thereof.

98. (new) The method according to claim 76, wherein the agronomic plant is selected from the group consisting of corn, cereals, barley, rye, rice, vegetables, clovers, legumes, beans, peas, alfalfa, sugar cane, sugar beets, tobacco, cotton, rapeseed (canola), sunflower, safflower, and sorghum.

99. (new) The method according to claim 76, wherein the agronomic plant is a member of the class Magnoliopsida.

100. (new) The method according to claim 76, wherein the agronomic plant is a member of the sub-family Papilionoideae or Faboideae.


101. (new) The method according to claim 76, wherein the agronomic plant is selected from the group consisting of *Pisum spp.*, *Medicago spp.*, *Arachis spp.*, *Glycine spp.*, *Vicia spp.*, *Vigna spp.*, trefoil, clovers and *Phaseolus spp.*

102. (new) The method according to claim 76, wherein the agronomic plant is a soybean plant.

103. (new) A method of increasing the vigor and/or the yield of an agronomic plant comprising treating the plant or its propagation material with a composition which comprises an effective amount of a fungicide which has no significant activity against fungal plant pathogens for such agronomic plant, wherein the plant or its propagation material possesses a transgenic event providing the plant with resistance to a herbicide and the treatment comprises foliar application of said herbicide.

104. (new) The method according to claim 103, wherein the herbicide is selected from the group consisting of glyphosate, glyphosinate, imidazilnone and STS system.

105. (new) The method according to claim 103, wherein the fungicide comprises a compound having the formula



wherein Z_1 and Z_2 are C or N and are part of an aromatic ring selected from benzene, pyridine, thiophene, furan, pyrrole, pyrazole, thiazole, benzothiophene and isothiazole;

A is selected from $--C(X)\text{-amine}$, $--C(O)\text{---}SR_3$, $--NH\text{---}C(X)R_4$, and $--C(=NR_3)\text{---}XR_7$;

B is $--W_m\text{---}Q(R_2)_3$ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R_4 ;

Q is C, Si, Ge, or Sn;

W is $--C(R_3)_p H_{(2-p)}--$; or when Q is C, W is selected from $--C(R_3)_p H_{(2-p)}--$, $--N(R_3)_m H_{(1-m)}--$, $--S(O)_p--$, and $--O--$;

X is O or S;

n is 0, 1, 2, or 3;

m is 0 or 1;

p is 0, 1, or 2;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;

b) C_1 - C_4 alkyl, alkenyl, alkynyl, C_3 - C_6 cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C_1 - C_4 alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxycarbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C_1 - C_4 alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C_1 - C_4 alkoxy, alkenoxy, alkynoxy, C_3 - C_6 cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo;

wherein two R groups may be combined to form a fused ring;

each R_2 is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R_4 or halogen; and wherein, when Q is C, R_2 may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino;

wherein two R_2 groups may be combined to form a cyclo group with Q;


R_3 is C_1 - C_4 alkyl;

R_4 is C_1 - C_4 alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino;

R_7 is C_1 - C_4 alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R_4 ; or an agronomic salt thereof.

106. (new) The method according to claim 103, wherein the fungicide is 4,5-dimethyl-*N*-2-propenyl-2-(trimethylsilyl)-3-thiophenecarboxamide.

107. (new) A method of increasing the vigor and/or the yield of an agronomic plant comprising treating the plant or its propagation material with a composition which comprises an effective amount of a fungicide which has no significant activity against fungal plant pathogens for such agronomic plant, wherein the treatment comprises treating the seed of the plant with an inoculant selected from the group consisting of *Azospirillum spp.*, *Rhizobium spp.*, *Bradyrhizobium spp.*, a mixture of *Rhizobium spp.* and *Bradyrhizobium spp.*, and a mixture of either *Rhizobium spp.*, or *Bradyrhizobium spp.* with any other microorganisms, and further includes foliar treatment of the plant with a compound having the formula



wherein Z_1 and Z_2 are C or N and are part of an aromatic ring selected from benzene, pyridine, thiophene, furan, pyrrole, pyrazole, thiazole, benzothiophene and isothiazole;

A is selected from $--C(X)\text{-amine}$, $--C(O)\text{---}SR_3$, $--NH\text{---}C(X)R_4$, and $--C(=NR_3)\text{---}XR_7$;

B is $--W_m\text{---}Q(R_2)_3$ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R_4 ;

Q is C, Si, Ge, or Sn;

W is $--C(R_3)_p\text{---}H_{(2-p)}\text{---}$; or when Q is C, W is selected from $--C(R_3)_p\text{---}H_{(2-p)}\text{---}$, $--N(R_3)_m\text{---}H_{(1-m)}\text{---}$, $--S(O)_p\text{---}$, and $--O\text{---}$;

X is O or S;

n is 0, 1, 2, or 3;

m is 0 or 1;

p is 0, 1, or 2;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;

b) C₁-C₄ alkyl, alkenyl, alkynyl, C₃-C₆ cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C₁-C₄ alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxy carbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C₁-C₄ alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C₁-C₄ alkoxy, alkenoxy, alkynoxy, C₃-C₆ cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl, alkylcarbonyloxy, alkoxy carbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo;

wherein two R groups may be combined to form a fused ring;

each R₂ is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R₄ or halogen; and wherein, when Q is C, R₂ may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino;

wherein two R₂ groups may be combined to form a cyclo group with Q;

R₃ is C₁-C₄ alkyl;

R₄ is C₁-C₄ alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino; and

R₇ is C₁-C₄ alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R₄; or an agronomic salt thereof.


108. (new) The method according to claim 107, wherein the plant or its propagation material possesses a transgenic event providing the plant with resistance to a herbicide selected from the group consisting of glyphosate, glyphosinate,

imidazilinone and STS system and the treatment further comprises foliar application of said herbicide.

109. (new) A method of increasing the vigor and/or the yield of an agronomic plant comprising treating the plant or its propagation material with a composition which comprises an effective amount of a fungicide which has no significant activity against fungal plant pathogens for such agronomic plant, wherein the step of treating the plant or its propagation material comprises applying the fungicide to the foliage of the plant, and wherein the plant or its propagation material possesses a transgenic event providing the plant with resistance to a herbicide and the step of applying the fungicide to the foliage of the plant comprises the application of the fungicide in combination with said herbicide.

110. (new) The method according to claim 109, wherein the herbicide is glyphosate.

111. (new) The method according to claim 109, wherein the fungicide comprises a compound having the formula



wherein Z_1 and Z_2 are C or N and are part of an aromatic ring selected from benzene, pyridine, thiophene, furan, pyrrole, pyrazole, thiazole, benzothiophene and isothiazole;

A is selected from $--C(X)\text{-amine}$, $--C(O)\text{---}SR_3$, $--NH\text{---}C(X)R_4$, and $--C(=NR_3)\text{---}XR_7$;

B is $--W_m\text{---}Q(R_2)_3$ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R_4 ;

Q is C, Si, Ge, or Sn;

W is $--C(R_3)_p H_{(2-p)}--$; or when Q is C, W is selected from $--C(R_3)_p H_{(2-p)}--$, $--N(R_3)_m H_{(1-m)}--$, $--S(O)_p--$, and $--O--$;

X is O or S;

n is 0, 1, 2, or 3;

m is 0 or 1;

p is 0, 1, or 2;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;

b) C_1 - C_4 alkyl, alkenyl, alkynyl, C_3 - C_6 cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C_1 - C_4 alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxycarbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C_1 - C_4 alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C_1 - C_4 alkoxy, alkenoxy, alkynoxy, C_3 - C_6 cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo;

wherein two R groups may be combined to form a fused ring;

each R_2 is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R_4 or halogen; and wherein, when Q is C, R_2 may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino;

wherein two R_2 groups may be combined to form a cyclo group with Q;

R_3 is C_1 - C_4 alkyl;

R_4 is C_1 - C_4 alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino;


R_7 is C_1 - C_4 alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R_4 ; or an agronomic salt thereof.

112. (new) The method according to claim 109, wherein the fungicide is 4,5-dimethyl-*N*-2-propenyl-2-(trimethylsilyl)-3-thiophenecarboxamide.

113. (new) A method of increasing the vigor and/or the yield of an agronomic plant except for wheat comprising treating an agronomic plant or its propagation material except for wheat with a composition comprising an effective amount of an active agent that has activity against *Gaeumannomyces graminis*, wherein the plant or its propagation material possesses a transgenic event providing the plant with resistance to a herbicide and the treatment comprises foliar application of said herbicide.

114. (new) The method according to claim 113, wherein the *Gaeumannomyces graminis* is of the variety *tritici*.

115. (new) The method according to claim 113, wherein the active agent comprises a compound having the formula



wherein Z_1 and Z_2 are C or N and are part of an aromatic ring selected from benzene, pyridine, thiophene, furan, pyrrole, pyrazole, thiazole, benzothiophene and isothiazole;

A is selected from $--C(X)-\text{amine}$, $--C(O)SR_3$, $--NH--C(X)R_4$, and $--C(=NR_3)--XR_7$;

B is $--W_m--Q(R_2)_3$ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R_4 ;

Q is C, Si, Ge, or Sn;

W is $--C(R_3)_p H_{(2-p)}--$; or when Q is C, W is selected from $--C(R_3)_p H_{(2-p)}--$, $--N(R_3)_m H_{(1-m)}--$, $--S(O)_p--$, and $--O--$;

X is O or S;

n is 0, 1, 2, or 3;

m is 0 or 1;

p is 0, 1, or 2;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;

b) C₁-C₄ alkyl, alkenyl, alkynyl, C₃-C₆ cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C₁-C₄ alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxycarbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C₁-C₄ alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C₁-C₄ alkoxy, alkenoxy, alkynoxy, C₃-C₆ cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo;

wherein two R groups may be combined to form a fused ring;

each R₂ is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R₄ or halogen; and wherein, when Q is C, R₂ may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino;

wherein two R₂ groups may be combined to form a cyclo group with Q;

R₃ is C₁-C₄ alkyl; and

R₄ is C₁-C₄ alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino;

R₇ is C₁-C₄ alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R₄ ;
or an agronomic salt thereof.

116. (new) The method according to claim 113, wherein the active agent is 4,5-dimethyl-*N*-2-propenyl-2-(trimethylsilyl)-3-thiophenecarboxamide.


117. (new) The method according to claim 113, where the treatment of the plant or its propagation material comprises treatment of a seed with an inoculant comprising *Azospirillum spp.*, or *Rhizobium spp.*, or *Bradyrhizobium spp.*, or a mixture of *Rhizobium spp.* and *Bradyrhizobium spp.*, or a mixture of either *Rhizobium spp.*, or *Bradyrhizobium spp.* with any other microorganisms.

118. (new) The method according to claim 113, wherein the herbicide is selected from the group consisting of glyphosate, glyphosinate, imidazilnone and STS system.

119. (new) A method of increasing the vigor and/or the yield of an agronomic plant except for wheat comprising treating an agronomic plant or its propagation material except for wheat with a composition comprising an effective amount of an active agent that has activity against *Gaeumannomyces graminis*, and treating the seed of the plant with an inoculant selected from the group consisting of *Azospirillum spp.*, *Rhizobium spp.*, *Bradyrhizobium spp.*, a mixture of *Rhizobium spp.* and *Bradyrhizobium spp.*, and a mixture of either *Rhizobium spp.*, or *Bradyrhizobium spp.* with any other microorganisms, and further includes foliar treatment of the plant with the active agent.

120. (new) The method according to claim 119, wherein the herbicide is glyphosate.

121. (new) The method according to claim 119, wherein the active agent comprises a compound having the formula



wherein Z_1 and Z_2 are C or N and are part of an aromatic ring selected from benzene, pyridine, thiophene, furan, pyrrole, pyrazole, thiazole, benzothiophene and isothiazole;

A is selected from --C(X)-amine, --C(O)—SR₃, --NH--C(X)R₄, and --C(=NR₃)--XR₇;

B is --W_m --Q(R₂)₃ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R₄;

Q is C, Si, Ge, or Sn;

W is --C(R₃)_p H_(2-p) --; or when Q is C, W is selected from --C(R₃)_p H_(2-p) --, --N(R₃)_m H_(1-m) --, --S(O)_p --, and --O--;

X is O or S;

n is 0, 1, 2, or 3;

m is 0 or 1;

p is 0, 1, or 2;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;

b) C₁-C₄ alkyl, alkenyl, alkynyl, C₃-C₆ cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C₁-C₄ alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxycarbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C₁-C₄ alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C₁-C₄ alkoxy, alkenoxy, alkynoxy, C₃-C₆ cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo;

wherein two R groups may be combined to form a fused ring;

each R₂ is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R₄ or halogen; and wherein, when Q is C, R₂ may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino;


wherein two R_2 groups may be combined to form a cyclo group with Q;
 R_3 is C_1 - C_4 alkyl;
 R_4 is C_1 - C_4 alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino;
 R_7 is C_1 - C_4 alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R_4 ;
 or an agronomic salt thereof.

122. (new) The method according to claim 119, wherein the active agent is 4,5-dimethyl-*N*-2-propenyl-2-(trimethylsilyl)-3-thiophenecarboxamide.

123. (new) A soybean plant or soybean seed which has been treated with a composition comprising an active agent which has activity against *Gaeumannomyces graminis* in an amount sufficient to increase the yield and/or the vigor of the soybean plant, wherein the seed is contacted with an inoculant comprising *Rhizobium spp.*, or *Bradyrhizobium spp.*, or a mixture of *Rhizobium spp.* and *Bradyrhizobium spp.*, or a mixture of either *Rhizobium spp.*, or *Bradyrhizobium spp.* with any other microorganisms.

124. (new) The soybean plant or soybean seed according to claim 123, wherein the *Gaeumannomyces graminis* is of the variety *tritici*.

125. (new) The soybean plant or soybean seed according to claim 123, wherein the active agent comprises a compound having the formula



wherein Z_1 and Z_2 are C or N and are part of an aromatic ring selected from benzene, pyridine, thiophene, furan, pyrrole, pyrazole, thiazole, benzothiophene and isothiazole;

A is selected from $--C(X)\text{-amine}$, $--C(O)\text{---}SR_3$, $--NH\text{---}C(X)R_4$, and $--C(=NR_3)\text{---}XR_7$;

B is $--W_m--Q(R_2)_3$ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R_4 ;

Q is C, Si, Ge, or Sn;

W is $--C(R_3)_p H_{(2-p)}--$; or when Q is C, W is selected from $--C(R_3)_p H_{(2-p)}--$, $--N(R_3)_m H_{(1-m)}--$, $--S(O)_p--$, and $--O--$;

X is O or S;

n is 0, 1, 2, or 3;

m is 0 or 1;

p is 0, 1, or 2;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;

b) C_1 - C_4 alkyl, alkenyl, alkynyl, C_3 - C_6 cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C_1 - C_4 alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxycarbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C_1 - C_4 alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C_1 - C_4 alkoxy, alkenoxy, alkynoxy, C_3 - C_6 cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo;

wherein two R groups may be combined to form a fused ring;

each R_2 is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R_4 or halogen; and wherein, when Q is C, R_2 may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino;

wherein two R_2 groups may be combined to form a cyclo group with Q;

R_3 is C_1 - C_4 alkyl;

R₄ is C₁-C₄ alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino;


R₇ is C₁-C₄ alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R₄; or an agronomic salt thereof.

126. (new) The soybean plant or soybean seed according to claim 123, wherein the active agent is 4,5-dimethyl-N-(2-propenyl)-2-(trimethylsilyl)-3-thiophenecarboxamide.

127. (new) The soybean plant or soybean seed according to claim 121, wherein the seed is contacted with a fungicide selected from the group consisting of fludioxonil, fluquinconazole, difenoconazole, captan, metalaxyl, carboxin and thiram.

128. (new) A plant or its propagation material of the family Fabaceae which has been treated with a composition comprising an active agent which has activity against *Gaeumannomyces graminis* in an amount sufficient to increase the yield and/or the vigor of said plant, wherein the plant or its propagation material has a transgenic event.

129. (new) The plant or its propagation material according to claim 128, wherein the active agent comprises a compound having the formula



wherein Z₁ and Z₂ are C or N and are part of an aromatic ring selected from benzene, pyridine, thiophene, furan, pyrrole, pyrazole, thiazole, benzothiophene and isothiazole;

A is selected from --C(X)-amine, --C(O)--SR₃, --NH--C(X)R₄, and --C(=NR₃)--XR₇;

B is --W_m--Q(R₂)₃ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R₄;

Q is C, Si, Ge, or Sn;

W is $--C(R_3)_p H_{(2-p)}--$; or when Q is C, W is selected from $--C(R_3)_p H_{(2-p)}--$, $--N(R_3)_m H_{(1-m)}--$, $--S(O)_p--$, and $--O--$;

X is O or S;

n is 0, 1, 2, or 3;

m is 0 or 1;

p is 0, 1, or 2;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;

b) C₁-C₄ alkyl, alkenyl, alkynyl, C₃-C₆ cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C₁-C₄ alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxycarbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C₁-C₄ alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C₁-C₄ alkoxy, alkenoxy, alkynoxy, C₃-C₆ cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo;

wherein two R groups may be combined to form a fused ring;

each R₂ is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R₄ or halogen; and wherein, when Q is C, R₂ may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino;

wherein two R₂ groups may be combined to form a cyclo group with Q;

R₃ is C₁-C₄ alkyl;


R₄ is C₁-C₄ alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino;

R₇ is C₁-C₄ alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R₄; or an agronomic salt thereof.

130. (new) The plant or its propagation material according to claim 128, wherein the active agent is 4,5-dimethyl-N-(2-propenyl)-2-(trimethylsilyl)-3-thiophenecarboxamide.

131. (new) A plant or its propagation material of the family Fabaceae which has been treated with a composition comprising an active agent which has activity against *Gaeumannomyces graminis* in an amount sufficient to increase the yield and/or the vigor of said plant, wherein the plant or its propagation material is the product of a QTL-based selective breeding program.

132. (new) The plant or its propagation material according to claim 131, wherein the active agent comprises a compound having the formula



wherein Z_1 and Z_2 are C or N and are part of an aromatic ring selected from benzene, pyridine, thiophene, furan, pyrrole, pyrazole, thiazole, benzothiophene and isothiazole;

A is selected from $--C(X)\text{-amine}$, $--C(O)\text{---}SR_3$, $--NH\text{---}C(X)R_4$, and $--C(=NR_3)\text{---}XR_7$;

B is $--W_m\text{---}Q(R_2)_3$ or selected from o-tolyl, 1-naphthyl, 2-naphthyl, and 9-phenanthryl, each optionally substituted with halogen or R_4 ;

Q is C, Si, Ge, or Sn;

W is $--C(R_3)_p\text{---}H_{(2-p)}\text{---}$; or when Q is C, W is selected from $--C(R_3)_p\text{---}H_{(2-p)}\text{---}$, $--N(R_3)_m\text{---}H_{(1-m)}\text{---}$, $--S(O)_p\text{---}$, and $--O\text{---}$;

X is O or S;

n is 0, 1, 2, or 3;

m is 0 or 1;

p is 0, 1, or 2;

each R is independently selected from

a) halo, formyl, cyano, amino, nitro, thiocyanato, isothiocyanato, trimethylsilyl, and hydroxy;

b) C₁-C₄ alkyl, alkenyl, alkynyl, C₃-C₆ cycloalkyl, and cycloalkenyl, each optionally substituted with halo, hydroxy, thio, amino, nitro, cyano, formyl, phenyl, C₁-C₄ alkoxy, alkylcarbonyl, alkylthio, alkylamino, dialkylamino, alkoxycarbonyl, (alkylthio)carbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulfinyl, or alkylsulfonyl;

c) phenyl, furyl, thienyl, pyrrolyl, each optionally substituted with halo, formyl, cyano, amino, nitro, C₁-C₄ alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, dialkylamino, haloalkyl, and haloalkenyl;

d) C₁-C₄ alkoxy, alkenoxy, alkynoxy, C₃-C₆ cycloalkyloxy, cycloalkenyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, dialkylamino, alkylcarbonylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, (alkylthio)carbonyl, phenylcarbonylamino, phenylamino, each optionally substituted with halo;

wherein two R groups may be combined to form a fused ring;

each R₂ is independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl and phenyl, each optionally substituted with R₄ or halogen; and wherein, when Q is C, R₂ may also be selected from halo, alkoxy, alkylthio, alkylamino, and dialkylamino;

wherein two R₂ groups may be combined to form a cyclo group with Q;

R₃ is C₁-C₄ alkyl;

R₄ is C₁-C₄ alkyl, haloalkyl, alkoxy, alkylthio, alkylamino, or dialkylamino;

R₇ is C₁-C₄ alkyl, haloalkyl, or phenyl, optionally substituted with halo, nitro, or R₄; or an agronomic salt thereof.

133. (new) The plant or its propagation material according to claim 131, wherein the active agent is 4,5-dimethyl-N-(2-propenyl)-2-(trimethylsilyl)-3-thiophenecarboxamide.